
UNCLASSIFIED

AD NUMBER AD834642 **NEW LIMITATION CHANGE** TO Approved for public release, distribution unlimited **FROM** Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; SEP 1966. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Releases Branch, Frederick, MD 21701. **AUTHORITY** Fort Detrick/SMUFD ltr dtd 15 Feb 1972

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

AD854642

TRANSLATION NO. 1844

DATE: 23 September, 1066

DD C

DDC AVAILABILITY NOTICE

Reproduction of this publication in whole or in part is prohibited. However, DDC is authorized to reproduce the publication for United States Government purposes.

STATEMENT NO UNCLASSIFIED

A the field mach

atti, 's Lich Releases Br

Marie of the first of the

DEPARTMENT OF THE ARMY Fort Detrick Frederick, Maryland

21701

3

USE OF IMMUNOFLUORESCENCE METHOD IN IMMUNOLOGICAL DIAGNOSIS (PRACTICAL APPLICATIONS)

La Presse Medicale (The Medical Press) Vol 71, No 55, 1963, pp 2740-2742 J. Thivolet,

A. Kratchko and

M. Sepetdjian

(Study accomplished in part with the assistance of the National Hygiene Institute.)

([The authors are affiliated with the] Hygiene Laboratory of the Medical Faculty of Lyon (Pr Sohier) and the Dermatology Service of the Antiquaille Hospital (Pr agr. Thivolet).)

Summery

Are the methods of immunofluorescence sufficiently used in immunological diagnosis? The search for antibodies by means of these methods appeared interesting in many bacterial diseases, especially syphilis, as well as diseases due to parasites, mushrooms, virus, and particularly the pneumopathy due to the Eaton agent, in diseases with auto-antibodies (lupus erythematosus, Hashimoto disease, myasthemia); but it seems that these methods have so far not yet been used as widely in practice as they should be.

Conclusion

The quack review of the applications of the immunofluorescence method in immunological diagnosis brings out the gap still existing between the rather numerous preliminary studies and their extension to broad practical use. On the basis of this latter criterion, we can eingle out 3 applications, the test applied to the diagnosis of syphilis (F.T.A. test); the diagnostic test for pneumopathies due to the Eston agent, and the test in which we look for antinuclear antibodies. We should really have standardised all of these tests by now and we might hope that other reactions more practical ones, at that, will be applied, at least as regards the

Easton agent.

But the current vogue of these techniques and their improvement lead us to believe that we might be able to add other studies, expanding the field of application of this method right now and confirming certain basic premises, to the currently probably incomplete list which we have established here.